



Relationship of preoperative neutrophil lymphocyte ratio with prognosis in gastrointestinal stromal tumors

Süleyman Kargin¹, Murat Çakır², Ebubekir Gündeş², Yusuf Yavuz², Hacı Hasan Esen³, M. Sinan İyisoğlu⁴, Naile Kökbudak³, Tevfik Küçükartallar²

ABSTRACT

Objective: Gastrointestinal stromal tumors are believed to play a role in immunosuppression with their local effect. In some cancers with associated inflammation, an increased pretreatment neutrophil lymphocyte ratio is considered as an indicator of poor prognosis. The aim of our study was to assess the relationship of increased blood neutrophil lymphocyte ratio with disease prognosis in patients with gastrointestinal stromal tumors.

Material and Methods: The data of 78 patients who underwent surgery with the diagnosis of primary gastrointestinal stromal tumor in our clinic were evaluated retrospectively. The preoperative neutrophil lymphocyte ratio in the peripheral blood was determined. The neutrophil lymphocyte ratio and its relationship with tumor risk and prognosis were compared. The data were evaluated by Pearson's correlation analysis and the Welch ANOVA test.

Results: The preoperative neutrophil lymphocyte ratio was significantly increased in the high-risk groups ($p < 0.05$). An increased preoperative neutrophil lymphocyte ratio was associated with shorter survival ($r = -0.32$, $p = 0.009$). In addition, an increase in the neutrophil lymphocyte ratio was associated with an increase in the mitotic activity of the tumor ($r = -0.364$, $p = 0.025$).

Conclusion: The preoperative neutrophil lymphocyte ratio in gastrointestinal stromal tumors can be used as an indicator of high-risk tumors and poor prognosis.

Keywords: Gastrointestinal stromal tumor, neutrophil lymphocyte ratio, prognosis

INTRODUCTION

Gastrointestinal stromal tumors (GIST) are recently identified, rare mesenchymal tumors of the gastrointestinal tract. Significant molecular changes that lead to GIST development have been defined over the last two decades (1, 2). With the introduction of tyrosine kinase inhibitor imatinib mesylate in clinic use, survival rates have significantly increased in even metastatic and high risk GIST (3-5). In a randomized phase III trial by Blank et al. (6), it was reported that more than 80% of GIST patients benefit from the tyrosine kinase inhibitor and that median survival in metastatic GIST was approximately 5 years (6). According to the criteria by the National Institute of Health (NIH) and the Armed Forces Institute of Pathology (AFIP), tumor size and mitotic activity are the most reliable and most widely used prognostic criteria in GISTs.

In recent years, the importance of cancer-related inflammation in determining survival and disease progression is becoming apparent for various cancer types (7, 8). Especially the patient's systemic inflammation status independently determines prognosis. However, the reason for this remains unclear. Systemic inflammatory response is particularly associated with changes in white blood cell count in peripheral blood, and particularly with neutrophilia along with relative lymphocytopenia (9). Based on these results, neutrophil-lymphocyte ratio (NLR) has been shown to be associated with poor prognosis in various types of cancer (10, 11). NLR is a feasible, reproducible and inexpensive indicator that can be used to detect systemic inflammation (12). Previous studies have reported a correlation between high neutrophil counts in peripheral blood and response to treatment and poor prognosis in patients receiving imatinib therapy due to unresectable GISTs (13, 14).

In this study, our aim was to investigate the correlation between preoperative peripheral blood NLR in primary GIST patients and prognosis.

MATERIAL AND METHODS

Patients

An ethical board approval was obtained from Necmettin Erbakan University Clinical Research Ethics Committee, No. 2012-69. Files of 78 patients who underwent surgery at Necmettin Erbakan University Meram Medical Faculty General Surgery Clinics between January 2006 and June 2012 and had a pathological diagnosis of GIST were retrospectively evaluated. Data on preoperative routine peripheral complete blood count, biochemistry parameters, coagulation measurements and pathology results were reviewed retrospectively. Patients who were operated on for emergent reasons were excluded from

¹Clinic of General Surgery, Konya Training and Research Hospital, Konya, Turkey

²Department of General Surgery, Necmettin Erbakan University Faculty of Medicine, Konya, Turkey

³Department of Medical Pathology, Necmettin Erbakan University Faculty of Medicine, Konya, Turkey

⁴Department of Medical Statistics, Necmettin Erbakan University Faculty of Medicine, Konya, Turkey

Address for Correspondence Süleyman Kargin

Konya Eğitim Araştırma Hastanesi, Genel Cerrahi Kliniği, Konya, Türkiye
Phone: +90 332 323 67 09
e-mail: drs.kargin@hotmail.com

Received: 27.07.2014


Accepted: 10.09.2014

©Copyright 2015

by Turkish Surgical Association

Available online at

www.uluscerrahidergisi.org



the study. The correlation between patients' age, gender, type of surgery performed, pathology, immunohistochemistry features, follow-up period and NLR were assessed.

Peripheral Blood Measurements

Last peripheral blood count measured before surgery was accepted as the preoperative value. It was ensured that the patient did not have any other reasons that may cause leukocytosis. Blood NLR values were obtained by dividing blood neutrophil count (neutrophil count/ μL) to lymphocyte count (lymphocyte count/ μL). Correlation between NLR and prognosis of patients was analyzed.

Statistical Analysis

Statistical Package for the Social Sciences 16.0 (SPSS Inc., Chicago, IL, USA) was used for statistical analysis. Binary correlations were assessed by Pearson correlation analysis. The comparison of the difference between groups was evaluated by ANOVA Welch test. Data were expressed by hazard ratio and 95% confidence intervals. $P < 0.05$ was considered to be statistically significant.

RESULTS

Demographic Data

Forty-one patients were female and 37 male, with a mean age of 68.8 (31-83). There were 35 gastric, 20 small intestine, 11 retroperitoneal, 5 colon, 2 rectum, 2 esophagus, and one duodenum, one appendix and one adrenal tumors (Table 1). 69 patients underwent curative surgery while 9 were considered as inoperable during surgery. In the patients with inoperable tumors, the organ involved were the stomach, small bowel and retroperitoneum, with 3 patients in each site. The criteria for inoperability were accepted as multiple liver metastases and peritoneal carcinomatosis. These patients were treated with chemotherapy.

When mitosis count of the tumors were analyzed in 50 high power fields, the mitotic activity was found to be ≤ 5 in 33, 6-10 in 18, and ≥ 10 in 27 patients. According to this data, there were 7 cases in the very low risk group, 11 in the low risk group, 18 in the moderate risk group and 42 in the high-risk group.

The mean follow-up period was 40 months (6-120 months). Recurrence was observed in eight patients. Six of these patients were re-operated. Cure was achieved in four of these patients, while two patients were considered inoperable and received chemotherapy. Two patients with recurrence received chemotherapy without surgery due to metastatic disease. Fourteen patients could not be reached after a mean follow-up of 1 year.

Prognostic Criteria and NLR

The median NLR was 3.5 (0.9 to 32.5). The evaluation of NLR according to pathologic risk groups showed that NLR level was significantly increased parallel to an increase in risk group ($p < 0.02$). According to Pearson correlation analysis, the NLR increase and decrease in the follow-up period was statistically significant ($r = -0.32$, $p = 0.009$, Figure 1). Similarly, there was a significant correlation between increased mitotic activity and NLR increase ($r = -0.364$, $p = 0.025$, Figure 1b). On the other hand, there was no correlation between NLR increase and tumor diameter and recurrence ($p = 0.189$, $r = 0.202$).

DISCUSSION

Gastrointestinal stromal tumors are rare mesenchymal neoplasms of the gastrointestinal tract. The recent implementation of targeted therapies based on advances in both biology and pathogenesis of GISTs has provided dramatic results. At present, there is no staging system for GIST. Pathologists believe that tumor size and mitotic index are the most reliable indicators of malignancy. Currently, the NIH tumor size and tumor risk category are used as independent prognostic criteria.

Since gastrointestinal stromal tumors are resistant to radiation therapy or chemotherapy, there was no effective treatment for metastatic disease or for tumors that could not be surgically removed until the discovery of tyrosine kinase inhibitors (15). Currently, the first choice treatment of recurrent or metastatic GIST is targeted therapy (imatinib). With this treatment, half of the patients respond to treatment, and disease progression stops in at least 75-85% (16). Multiple tyrosine kinase inhibitor sunitinib is used in patients with imatinib resistance (17). Our patients high-risk patients according to AFIP criteria with inoperable disease were treated with targeted therapy. Six patients with an unresectable lesion showed partial remission with imatinib 400 mg / day treatment for 12 months. Curative

surgery was performed in the re-operation after down staging. The treatment of three patients is still on-going.

The role of immune response in the natural history of gastrointestinal stromal tumors is becoming increasingly evident. Inflammation refers to the presence of immune cells in the tumor, which is clinically associated with poor prognosis. Indeed, Balachandran et al. (14) has shown that intratumoral T-cell response is induced in GISTs. However, the CD4 / CD8 ratio is relatively deteriorated, and tumor cells are suppressed by indoleamine 2,3-dioxygenase, which is an immunosuppressive enzyme. The lack of general immune response due to impairment of T cell function may decrease tumor-specific response. This mechanism is important in imatinib sensitivity. Another mechanism is stimulation of tumor cell growth and angiogenesis by inflammatory cells, and induction of metastasis through mediators they secrete (18).

Preoperative prediction of prognosis in gastrointestinal stromal tumors is not only important in providing data on postoperative survival for clinicians but also in determining adjuvant treatment. Today, adjuvant imatinib therapy is given to high-

risk patients according to the NIH criteria. Patients with three selected risk factors; tumor size, tumor location, and high mitotic index have high recurrence rates and benefit greatly from adjuvant therapy (19). However, these criteria can be only determined by postoperative pathology evaluation. Parameters that may inform clinicians on tumor characteristics preoperatively on non-metastatic GIST are limited. In particular, we can only make estimation on tumor characteristics preoperatively based on tumor size according to the AFIP and the NIH criteria. We aimed to identify a new criterion that is both efficient and easily applicable in determining the character of GIST tumors preoperatively. We considered that in this way the surgeon will be informed on tumor characteristics before the operation and may help in determining the patient's treatment plan.

NLR is one of the hematological markers of systemic inflammatory response, and is being used as a reliable method both as an indicator of inflammation and in determining survival in several types of cancer. Öztürk et al. (20) have reported that NLR significantly increased in young patients with acute coronary syndrome. Kapçı et al. (21) and Kahramanca et al. (22) emphasized the importance of NLR increase in the diagnosis of acute appendicitis. The prognostic significance of NLR in various types of cancer has been investigated in numerous studies (Table 2) (23-31). Presence of systemic inflammation reaction is used as a prognostic marker for various types of cancer (27, 30). In our study, we determined that preoperative NLR is a strong predictor of both mitotic activity and pathologic tumor risk for GIST. In addition, we identified that enhanced NLR is associated with poor prognosis. This finding indicates that the NLR value, which is an applicable and cheap preoperative method, can be used as an independent predictor of prognosis in GIST.

CONCLUSION

Preoperative NLR may be used as a feasible, inexpensive and reliable prognostic factor that both provides information on the tumor risk and helps determine prognosis in gastrointestinal stromal tumors.

Ethics Committee Approval: Ethics committee approval was received for this study from the ethics committee of Necmettin Erbakan University Clinical Research (2012-69).

Informed Consent: Patient consent form was not obtained from patients since this study is a database search.

Peer-review: Externally peer-reviewed.

Table 1. Distribution of patients according to organ and type of surgery

Organ location	n	Operation
Stomach	16	Gastric wedge resection
	7	Distal subtotal gastrectomy
	7	Total gastrectomy
	2	Gastric wedge resection + segmentectomy
Small bowel	17	Partial bowel resection
Colon	5	Left hemicolectomy
Rectum	2	Low anterior resection
Oesophagus	2	Distal oesophagectomy
Duodenum	1	Whipple's procedure
Appendix	1	Appendectomy
Adrenal gland	1	Adrenalectomy
Retroperitoneum	8	Excision
Inoperable patients	9	Laparotomy + biopsy

Table 2. Evaluation of correlation between NLR and recurrence and survival after curative surgery in various cancer types

Study	Center	Organ	n	p Value	Threshold value	Footnote
Wang (23)	Guangdong (China)	Liver	101	<0.001	>3	High NLR is independently associated with disease free survival
Halazun (24)	New York (USA)	Liver	150	0.005	>5	High NLR is associated with recurrence and mortality rate
Bertuzzo (25)	Bologna (Italy)	Liver	219	<0.001	>5	High NLR is associated with low survival and negatively effects disease-free survival. NLR is an independent prognostic factor
Sato (26)	Shizuoka (Japan)	Oesophagus	83	0.043	>2.2	NLR is associated with pathologic response to treatment
Miyata (27)	Osaka (Japan)	Oesophagus	152	NS	>4	High NLR is associated with survival but not an independent prognostic factor
Sharaiha (28)	New York (USA)	Oesophagus	339	<0.001	>5	High NLR is associated with disease-free survival and effects survival independent of tumor type

NLR: neutrophil-lymphocyte ratio, NS: not significant

Author Contributions: Concept - S.K., T.K.; Design - S.K.; Supervision - M.Ç., Y.Y.; Funding - Y.Y., E.G.; Materials - S.K.; Data Collection and/or Processing - S.K., Y.Y.; Analysis and/or Interpretation - H.H.E., N.K.; Literature Review - M.S.İ.; Writer - S.K.; Critical Review - M.S.İ.

Conflict of Interest: No conflict of interest was declared by the authors.

Financial Disclosure: The authors declared that this study has received no financial support.

REFERENCES

- Hirota S, Isozaki K, Moriyama Y, Hashimoto K, Nishida T, Ishiguro S, et al. Gain-of-function mutations of c-kit in human gastrointestinal stromal tumors. *Science* 1998; 279: 577-580. [\[CrossRef\]](#)
- Heinrich MC, Corless CL, Duensing A, McGreevey L, Chen CJ, Joseph N, et al. PDGFRA activating mutations in gastrointestinal stromal tumors. *Science* 2003; 299: 708-710. [\[CrossRef\]](#)
- Joensuu H, Roberts PJ, Sarlomo-Rikala M, Andersson LC, Tervahartiala P, Tuveson D, et al. Effect of the tyrosine kinase inhibitor STI571 in a patient with a metastatic gastrointestinal stromal tumor. *N Engl J Med* 2001; 344: 1052-1056. [\[CrossRef\]](#)
- Dematteo RP, Ballman KV, Antonescu CR, Maki RG, Pisters PW, Demetri GD, et al. Adjuvant imatinib mesylate after resection of localised, primary gastrointestinal stromal tumour: a randomised, double-blind, placebo-controlled trial. *Lancet* 2009; 373: 1097-1104. [\[CrossRef\]](#)
- Fletcher CD, Berman JJ, Corless C, Gorstein F, Lasota J, Longley BJ, et al. Diagnosis of gastrointestinal stromal tumors: A consensus approach. *Hum Pathol* 2002; 33: 459-465. [\[CrossRef\]](#)
- Blanke CD, Rankin C, Demetri GD, Ryan CW, Von Mehren M, Benjamin RS, et al. Phase III randomized, intergroup trial assessing imatinib mesylate at two dose levels in patients with unresectable or metastatic gastrointestinal stromal tumors expressing the kit receptor tyrosine kinase: S0033. *J Clin Oncol* 2008; 26: 626-632. [\[CrossRef\]](#)
- Hanahan D, Weinberg RA. The hallmarks of cancer. *Cell* 2000; 100: 57-70. [\[CrossRef\]](#)
- Hanahan D, Weinberg RA. Hallmarks of cancer: the next generation. *Cell* 2011; 144: 646-674. [\[CrossRef\]](#)
- Zahorec R. Ratio of neutrophil to lymphocyte counts-rapid and simple parameter of systemic inflammation and stress in critically ill. *Bratisl Lek Listy* 2001; 102: 5-14.
- Yamanaka T, Matsumoto S, Teramukai S, Ishiwata R, Nagai Y, Fukushima M. The baseline ratio of neutrophils to lymphocytes is associated with patient prognosis in advanced gastric cancer. *Oncology* 2007; 73: 215-220. [\[CrossRef\]](#)
- Cho H, Hur HW, Kim SW, Kim SH, Kim JH, Kim YT, et al. Pre-treatment neutrophil to lymphocyte ratio is elevated in epithelial ovarian cancer and predicts survival after treatment. *Cancer Immunol Immunother* 2009; 58: 15-23. [\[CrossRef\]](#)
- Perez DR, Baser RE, Cavnar MJ, Balachandran VP, Antonescu CR, Tap WD, et al. Blood neutrophil-to-lymphocyte ratio is prognostic in gastrointestinal stromal tumor. *Ann Surg Oncol* 2013; 20: 593-599. [\[CrossRef\]](#)
- Van Glabbeke M, Verweij J, Casali PG, Le Cesne A, Hohenberger P, Ray-Coquard I, et al. Initial and late resistance to imatinib in advanced gastrointestinal stromal tumors are predicted by different prognostic factors: a European Organisation for Research and Treatment of Cancer-Italian Sarcoma Group-Australasian Gastrointestinal Trials Group study. *J Clin Oncol* 2005; 23: 5795-5804. [\[CrossRef\]](#)
- Gastrointestinal Stromal Tumor Meta-Analysis Group (Meta-GIST). Comparison of two doses of imatinib for the treatment of unresectable or metastatic gastrointestinal stromal tumors: a meta-analysis of 1,640 patients. *J Clin Oncol* 2010; 28: 1247-1253. [\[CrossRef\]](#)
- Ji F, Wang ZW, Wang LJ, Ning JW, Xu GQ. Clinicopathological characteristics of gastrointestinal mesenchymal tumors and diagnostic value of endoscopic ultrasonography. *J Gastroenterol Hepatol* 2008; 23: 318-324. [\[CrossRef\]](#)
- Verweij PJ, Casali PG, Zalcberg PJ, LeCesne A, Reichardt P, Blay JY, et al. Progression-free survival in gastrointestinal stromal tumours with high-dose imatinib: randomized trial. *Lancet* 2004; 364: 1127-1134. [\[CrossRef\]](#)
- Rock EP, Goodman V, Jiang JX, Mahjoob K, Verbois SL, Morse D, et al. Food and drug administration drug approval summary: Sunitinib malate for the treatment of gastrointestinal stromal tumor and advanced renal cell carcinoma. *Oncologist* 2007; 12: 107-113. [\[CrossRef\]](#)
- Azuma T, Matayoshi Y, Odani K, Sato Y, Sato Y, Nagase Y, et al. Preoperative neutrophil-lymphocyte ratio as an independent prognostic marker for patients with upper urinary tract urothelial carcinoma. *Clin Genitourin Cancer* 2013; 11: 337-341. [\[CrossRef\]](#)
- Tanimine N, Tanabe K, Suzuki T, Tokumoto N, Ohdan H. Prognostic criteria in patients with gastrointestinal stromal tumors: a single center experience retrospective analysis. *World J Surg Oncol* 2012; 10: 43. [\[CrossRef\]](#)
- Öztürk S, Erdem A, Özlü MF, Ayhan S, Erdem K, Özayaşar M, et al. Assessment of the neutrophil to lymphocyte ratio in young patients with acute coronary syndromes. *Arch Turk Soc Cardiol* 2013; 41: 284-289. [\[CrossRef\]](#)
- Kapçı M, Türkdöğän KA, Duman A, Avcil M, Gülen B, Uğurlu Y, et al. Biomarkers in the diagnosis of acute appendicitis. *J Clin Exp Invest* 2014; 5: 250-255. [\[CrossRef\]](#)
- Kahramanca S, Ozgehan G, Seker D, Gökce El, Seker G, Tunç G, et al. Neutrophil-to-lymphocyte ratio as a predictor of acute appendicitis. *Ulus Travma Acil Cerrahi Derg* 2014; 20: 19-22. [\[CrossRef\]](#)
- Jablonska J, Leschner S, Westphal K, Lienenklaus S, Weiss S. Neutrophils responsive to endogenous IFN-beta regulate tumor angiogenesis and growth in a mouse tumor model. *J Clin Invest* 2010; 120: 1151-1164. [\[CrossRef\]](#)
- Müller I, Munder M, Kropf P, Hänsch GM. Polymorphonuclear neutrophils and T lymphocytes: strange bedfellows or brothers in arms? *Trends Immunol* 2009; 30: 522-530. [\[CrossRef\]](#)
- Guthrie GJ, Charles KA, Roxburgh CS, Horgan PG, McMillan DC, Clarke SJ. The systemic inflammation-based neutrophil lymphocyte ratio: experience in patients with cancer. *Crit Rev Oncol Hematol* 2013; 88: 218-230. [\[CrossRef\]](#)
- Wang GY, Yang Y, Li H, Zhang J, Jiang N, Li MR, et al. A scoring model based on neutrophil to lymphocyte ratio predicts recurrence of HBV-associated hepatocellular carcinoma after liver transplantation. *PLoS One* 2011; 6: e25295. [\[CrossRef\]](#)
- Halazun KJ, Hardy KJ, Rana AA, Woodland DC, Luyten EJ, Mahadev S, et al. Negative impact of neutrophil-lymphocyte ratio on outcome after liver transplantation for hepatocellular carcinoma. *Ann Surg* 2009; 250: 141-151. [\[CrossRef\]](#)
- Bertuzzo VR, Cescon M, Ravaoli M, Grazi GL, Ercolani G, Del Gaudio M, et al. Analysis of factors affecting recurrence of hepatocellular carcinoma after liver transplantation with a special focus on inflammation markers. *Transplantation* 2011; 91: 1279-1285. [\[CrossRef\]](#)
- Sato H, Tsubosa Y, Kawano T. Correlation between the pretherapeutic neutrophil to lymphocyte ratio and the pathologic response to neoadjuvant chemotherapy in patients with advanced esophageal cancer. *World J Surg* 2012; 36: 617-622. [\[CrossRef\]](#)
- Miyata H, Yamasaki M, Kurokawa Y, Takiguchi S, Nakajima K, Fujiwara Y, et al. Prognostic value of an inflammation-based score in patients undergoing pre-operative chemotherapy followed by surgery for esophageal cancer. *Exp Ther Med* 2011; 2: 879-885. [\[CrossRef\]](#)
- Sharaiha RZ, Halazun KJ, Mirza F, Port JL, Lee PC, Neugut AI, et al. Elevated preoperative neutrophil:lymphocyte ratio as a predictor of postoperative disease recurrence in esophageal cancer. *Ann Surg Oncol* 2011; 18: 3362-3369. [\[CrossRef\]](#)