



# Femoral artery aneurysm developed on intimal sarcoma: Case report

Fatih Yanar<sup>1</sup> , Gizem Öner<sup>2</sup> , Sercan Yüksel<sup>1</sup> , Ahmet Salduz<sup>3</sup> , İbrahim Fethi Azamat<sup>1</sup> , Beyza Özçınar<sup>1</sup> , Bilge Bilgic<sup>4</sup> , Rana Günöz Cömert<sup>5</sup> , Erol Kozanoğlu<sup>6</sup> , Mehmet Kurtoğlu<sup>1</sup> 

<sup>1</sup> Department of General Surgery, İstanbul University Faculty of Medicine, İstanbul, Türkiye

<sup>2</sup> Multidisciplinary Oncologic Centre (MOCA), Antwerpen University Hospital, Antwerpen, Belgium

<sup>3</sup> Department of Orthopaedic Surgery and Traumatology, İstanbul University Faculty of Medicine, İstanbul, Türkiye

<sup>4</sup> Department of Pathology, İstanbul University Faculty of Medicine, İstanbul, Türkiye

<sup>5</sup> Department of Radiology, İstanbul University Faculty of Medicine, İstanbul, Türkiye

<sup>6</sup> Department of Plastic and Reconstructive Surgery, İstanbul University Faculty of Medicine, İstanbul, Türkiye

## ABSTRACT

Intimal angiosarcoma is a rare vascular malignancy, and diagnosis is very difficult due to nonspecific symptoms. There are controversial points regarding the diagnosis, treatment and follow-up of intimal angiosarcomas. The purpose of this case report was to evaluate the diagnosis and treatment process of a patient diagnosed with femoral artery intimal angiosarcoma. Furthermore, in line with previous studies, it was aimed to illuminate controversial points. A 33-year-old male patient, who had been operated on due to ruptured femoral artery aneurysm, was diagnosed with intimal angiosarcoma with the pathology result. Recurrence was observed during clinical follow-up, and the patient was treated with chemotherapy and radiotherapy. Since there was no response to treatment, the patient underwent aggressive surgery including the surrounding tissues. No recurrence or metastasis was observed in the patient's 10<sup>th</sup> month follow-up. Although intimal angiosarcoma is rare, it should be considered in differential diagnosis when femoral artery aneurysm is detected. The most important step in treatment is aggressive surgery, but adding chemo-radiotherapy to the treatment should be considered.

**Keywords:** Intimal angiosarcoma, surgery, chemo-radiotherapy

## INTRODUCTION

Vascular malignancies are observed rarely, and angiosarcoma constitutes less than one percent of them, consisting of two subtypes: intimal type and mural type (1-3). Intimal angiosarcoma originates from the intimal layer of the arterial wall. It can develop at any level of the aorta and manifests different symptoms depending on its levels. Symptoms usually emerge as a result of stenosis, obstruction or thromboembolic events (3-5). It is difficult to diagnose intimal angiosarcoma due to nonspecific clinical presentation. In addition, intimal sarcoma diagnosed with femoral artery aneurysm is a very rare case. Surgery with negative tumour margins is a cornerstone of the intimal angiosarcoma treatment (6-7). It becomes apparent that chemotherapy and radiotherapy play a role in sarcoma treatment, but less attention has been paid to specific angiosarcoma treatment (8-13).

This case report attempted to shed light on how an intimal angiosarcoma case should be handled and addressed a number of dilemmas regarding diagnosis, treatment options and follow-up methods.

## CASE REPORT

A 33-year-old male presented with a complaint of swelling and pain in the right groin for a month. His medical history and family history were not remarkable. On physical examination, the pulses of bilateral lower extremities were palpable, and the ankle arm pressure index was measured as 1.2 on the right and 1.3 on the left. Arterial Doppler ultrasound and Computed Tomography (CT) imaging revealed a ruptured right femoral artery aneurysm that was located 5.5 x 4.5 cm periphery of the right main femoral artery and was measured 7 mm at its thickest site (Figure 1).

**Cite this article as:** Yanar F, Öner G, Yüksel S, Salduz A, Azamat İF, Özçınar B, et al. Femoral artery aneurysm developed on intimal sarcoma: Case report. Turk J Surg 2022; 38 (3): 306-311.

### Corresponding Author

Gizem Öner

E-mail: onergizem@hotmail.com


Received: 11.10.2020

Accepted: 05.05.2022

Available Online Date: 19.09.2022

© Copyright 2022 by Turkish Surgical Society Available online at www.turkjsurg.com

DOI: 10.47717/turkjsurg.2022.4381




**Figure 1.** **A.** In the arterial phase extremity CT angiography, pseudo-aneurysm is observed in the right common femoral artery with an irregular wall in the right common femoral artery in the arterial phase at the level of the femoral neck. **B.** In the coronal CT reformat maximum intensity projection image, the contours and extension of the pseudo-aneurysm in the common femoral artery. **C.** In 3-D CT reformat image, the relationship between pseudo-aneurysm, neck and common femoral artery can be observed.

The abdominal aorta, main iliac and external iliac arteries were detected normal. The patient was investigated for systemic autoimmune diseases and vasculitis, and no remarkable result was obtained. In addition, the patient had no history of trauma.


The patient underwent aneurysmectomy, and common femoral artery reconstruction was performed with a synthetic ringed polytetrafluoroethylene (PTFE) graft by preserving the external iliac and deep femoral arteries. After the operation, distal pulses of the patient were palpable and the ankle arm pressure index was measured as 1.2 on the right. Pathological examination of the specimen revealed pleomorphic-undifferentiated sarcoma originating from the intima with Vimentin (+) and a margin greater than 1 cm (Figure 2). Thoracic CT and Positron Emission Tomography (PET) showed no distant metastasis. The patient was evaluated by the oncology unit after the surgery, and clinical follow-up was decided due to tumour free margin and no metastasis. In the third month of follow-up, the patient complained about swelling and pain in the right groin. Femoral artery pulses could not be measured on the right lower extremity. Magnetic Resonance Imaging (MRI) and PET imaging showed a relapse of sarcoma that was measured as 9 x 7 cm. Recurrent sarcoma was found to extend the iliac muscle at the level of the right iliac crest and encircled the deep and superficial femoral arteries 360 degrees. In addition, a 37 x 27 mm metastatic lesion at the level of the collum femoris and trochanter minor

was detected (Figure 3). Chemo-radiotherapy was decided in order to perform surgical resection with a wider tumour negative margin.

Two cycles of chemotherapy at three-week intervals (150 mg Adriablastin + 6000 mg ifosfamide plus + 3000 mg mesna) and 28 Gy/8 fr radiotherapy were administered to the patient. During chemo-radiotherapy, deep vein thrombosis developed within the right femoral vein and subcutaneous enoxaparin sodium 2 x 8000 IU was initiated at the treatment dose. Surgery was decided since the MRI revealed no regression in the relapse lesion and metastatic lesions after chemotherapy, and surgical resection was considered possible in preoperative evaluation. Eight months after the first operation, the skin including the previous incision line, subcutaneous tissue, proximal femur and neighbouring muscles, external iliac artery and vein, and superficial femoral artery and vein were removed en bloc, and then vein ligation was performed (Figure 4). A femur prosthesis was placed for reconstruction and by-pass was performed with a synthetic ringed PTFE graft at the level of bifurcation between the main iliac artery proximally and superficial femoral artery distally. The femoral vein and main iliac vein were ligated (Figure 5). Plastic and reconstructive surgery was also included in the reconstruction part of the operation. A vertical pedicular rectus abdominis flap was removed to the left side of the deep inferior epigastric artery. The flap was transposed to the




**Figure 2.** Thrombus in the vessel lumen and endovascular tumour with intimal component. **A.** Hematoxylin and eosin (HE), x100, **B.** HE, x200, **C.** Vimentin (+), x100, **D.** Pleomorphic malignant tumour cells (HE, x200).




**Figure 3. A.** In contrast-enhanced MRI scan, in the axial MRI view of T1A sequence post contract fat suppression, common femoral artery (short yellow arrow), the central cystic-necrotic infiltrative mass lesion with dense heterogeneous contrast involvement in the pseudo-aneurysm region (long yellow arrow), metastatic lesion is observed in the medulla-trabecular area with intense contrast enhancement (orange arrow). **B.** In contrast-enhanced MRI scan, in the sagittal MRI view of T1A sequence post contract fat suppression, infiltrative, heterogeneous contrasting mass lesion and femoral head metastatic lesion (orange arrow). **C.** In the arterial phase extremity CT angiography, after endovascular stent graft inserted in the common femoral artery, the stent lumen open and the pseudo-aneurysm sac with thrombosis.

oncologic resection area. The flap of the left rectus abdominal muscle, skin and subcutaneous tissue, anterior region of the femoral prosthesis and the resection region were closed primarily (Figure 6).

Postoperatively, all pulses of the right lower extremity were palpable, and the ankle arm pressure index was measured as 1. In postoperative follow-up, flap circulation was good. Postoperatively, oral acetylsalicylic acid 100 mg, subcutaneous enoxaparin sodium 4000 IU and intermittent pneumatic compression




**Figure 4.** **A.** The clinical appearance of the wide resection specimen with the proximal part of the femur. **B.** The clinical appearance of the wide resection specimen with the proximal part of the femur.



**Figure 5.** Postoperative picture of the resection area. Synthetic ringed polytetrafluoroethylene (PTFE) graft was placed between iliac artery and superficial femoral artery. The femoral vein and main iliac vein were ligated.

on bilateral lower extremities were initiated for DVT prophylaxis. Pathological examination reported an undifferentiated high grade sarcoma with 85% necrosis and negative surgical margins in vascular, soft tissue, bone tissue structures. The patient was discharged on postoperative day 10 with oral acetylsalicylic acid 100 mg/day. On control, distal pulses of the right lower extremity were palpable and ankle arm pressure index was 1.1.



**Figure 6.** The flap of left rectus abdominis muscle, skin and subcutaneous tissue, anterior region of femoral prosthesis and resection region were closed primarily.

Two months after the second operation, the patient received paclitaxel 150 mg/week treatment for 12 weeks. No local relapse and distant metastasis were detected in the follow up control visit performed in the tenth month.

#### DISCUSSION

Intimal sarcomas are quite rare undifferentiated pleomorphic sarcomas originating from the tunica intima (1-3). According to the results of a review by Sebenik et al., 90% of intimal sarcomas originate from the aorta and 10% of them originate from the iliac and femoral arteries (4). Clinical findings vary depending

on the level of the affected vessel, whether there is stenosis in the aortic lumen or not, or whether thromboembolic metastasis developed or not. Intimal sarcomas form polypoid masses growing toward the lumen, which causes more commonly obstruction or thrombo-emboli symptoms (1-5). Diagnosis of intimal angiosarcoma is difficult due to non-specific clinic presentation and extreme rarity. Aortic angiosarcoma presents as an asymptomatic thoracic aneurysm, dyspnea, sudden death or distal embolization. However, the case of intimal sarcoma diagnosed with femoral artery aneurysm and/or arterial rupture is very rare in the literature. The present case was admitted to our clinic with non-specific complaints and was primarily investigated for vasculitis. Diagnostic approach for intimal angiosarcoma is based on a high degree of suspicion. An alternative explanation is that in the differential diagnosis of an aneurysm, which is not associated with vasculitis and traumatic status, intimal sarcomas should be considered.

While planning the treatment of intimal sarcoma, MRI is important for the assessment of local invasion and determining resectability whereas intravenous contrast CT is important in assessing distant metastasis. There are insufficient data on the use of PET scans in this type of tumour (5,6,13). Although there is no defined algorithm for the treatment of intimal sarcoma, aggressive surgical resection including neighbouring tissues is recommended in non-metastatic cases. Obtaining negative surgical margin is the most important factor for the prevention of relapse, which is fostered by some local recurrence studies (6,7). From this point of view, it appears that frozen section (FS) examination is also recommended for the assessment of resection intra-operatively. In the first operation, FS was not performed because it was not evaluated as intimal sarcoma. Diagnosis of intimal sarcoma was made after surgery, and no additional surgical intervention was considered since the surgical margin was greater than 1 cm. In a study conducted by Alamanda et al., positive surgical margin has been shown to increase relapse rates by 12-38% over a six-year follow-up (7). Re-resection should be performed, if possible, in patients with surgical margin positivity. Administration of radiotherapy is recommended if re-resection is not possible (8,9).

The efficacy of radiotherapy and chemotherapy in case of intimal sarcomas is contradictory. However, there are studies indicating that chemotherapy and radiotherapy administered after surgery have positive effects on survival (10,11). In addition, recent literature on extremity soft tissue sarcoma stresses that preoperative chemo-radiation improved overall survival and disease-free survival, but it is not a standard treatment (10). Beane JD et al. have shown that postoperative RT improves local control in high-grade lesions such as intimal sarcoma tumours, but its effect on overall survival has not been demonstrated (12). On the other hand, previous research has mostly

tended to focus on soft tissue sarcoma rather than on intimal angiosarcoma. In our case, it was decided to continue with adjuvant chemotherapy and radiotherapy after early recurrence, but a satisfactory response could not be obtained. Therefore, en bloc resection including bone, soft tissue and vessels was performed.

There is limited data available in the literature on surveillance strategies for intimal angiosarcoma. There is no exact data about which imaging method is better for follow-up. During the first-year follow-up, MRI for local recurrence and CT for distant metastasis may be preferred every four to six months. In this case, no evidence of relapse was found in the 10-month follow-up period.

## CONCLUSION

We presented a rare femoral angiosarcoma case, which was treated with chemo-radiotherapy and was operated aggressively to achieve tumour negative margins. When femoral artery aneurysm is detected in patients with non-specific symptoms, intimal sarcoma should be considered in differential diagnosis. The most important step in treatment is aggressive surgery, but adding chemo-radiotherapy to the treatment should be considered.

**Peer-review:** Externally peer-reviewed.

**Author Contributions:** Concept – F.Y., G.Ö., S.Y., B.Ö., M.K.; Design – F.Y., G.Ö.; Supervision – A.S., İ.F.A., B.Ö.; Materials – F.Y., G.O., S.Y., B.B., R.G.C., E.K.; Data Collection and/or Processing – F.Y., G.Ö., A.S., İ.F.A., S.Y.; Literature Search – F.Y., G.Ö.; Writing Manuscript – F.Y., G.O., S.Y., A.S., İ.F.A., B.O., B.B., R.G.C., E.K., M.K.; Critical Reviews – F.Y., G.Ö., S.Y., A.S., İ.F.A., B.Ö., M.K.

**Conflict of Interest:** The authors have no conflicts of interest to declare.

**Financial Disclosure:** The authors declared that this study has received no financial support.

## REFERENCES

1. Pucci A, De Martino A, Levantino M, Berchiolli R, Basolo F, Bortolotti U. Intimal sarcoma of the descending aorta mimicking aortitis. *Aorta* 2016; 4: 142-5. <https://doi.org/10.12945/j.aorta.2016.16.022>
2. Anthonissen N, Menting T, Verkroost M, Morshuis W. Angiosarcoma of the descending aorta, diagnostic difficulties. *EJVES Short Rep* 2016; 32: 4-6. <https://doi.org/10.1016/j.ejvssr.2016.04.002>
3. Ozbek C, Emreca B, Calli AO, Gurbuz A. Intimal sarcoma of the pulmonary artery with retrograde extension into the pulmonic valve and right ventricle, *Texas Hear Inst J* 2007; 34: 119-21.
4. Sebenik M, Ricci AJR, DiPasquale B, Mody K, Pytel P, Jee KJ. et al. Undifferentiated intimal sarcoma of large systemic blood vessels: Report of 14 cases with immunohistochemical profile and review of the literature. *Am J Surg Pathol* 2005; 29:1184-93. <https://doi.org/10.1097/01.pas.0000159774.70288.7d>
5. Ebaugh JL, Yuan M, Hu J, Chen A, Raffetto JD. Intimal sarcoma of the superficial femoral artery with osteosarcomatous differentiation, *J Vasc Surg* 2011; 53: 1394-97. <https://doi.org/10.1016/j.jvs.2010.11.098>

6. Staats P, Tavora F, Burke AP. Intimal sarcomas of the aorta and iliofemoral arteries: A clinicopathological study of 26 cases, *Pathology* 2014; 46: 596-603. <https://doi.org/10.1097/PAT.0000000000000182>
7. Alamanda VK, Crosby SN, Archer KR, Song Y, Schwartz HS, Holt GE. Predictors and clinical significance of local recurrence in extremity soft tissue sarcoma. *Acta Oncol* 2013; 52: 793-802. <https://doi.org/10.3109/0284186X.2012.711953>
8. Arai E, Nishida Y, Tsukushi S, Wasa J, Ishiguro N. Clinical and treatment outcomes of planned and unplanned excisions of soft tissue sarcomas. *Clin Orthop Relat Res* 2010; 468: 3028-34. <https://doi.org/10.1007/s11999-010-1392-7>
9. Cahlon O, Spierer M, Brennan MF, Singer S, Alektiar KM. Long-term outcomes in extremity soft tissue sarcoma after a pathologically negative re-resection and without radiotherapy. *Cancer* 2008; 112: 2774-79. <https://doi.org/10.1002/cncr.23493>
10. Mullen JT, Kobayashi W, Wang JJ, Harmon DC, Choy E, Hornicek FJ, et al. Long-term follow-up of patients treated with neoadjuvant chemotherapy and radiotherapy for large, extremity soft tissue sarcomas. *Cancer* 2012; 118: 3758-65. <https://doi.org/10.1002/cncr.26696>
11. Grobmyer SR, Maki RG, Demetri GD, Mazumdar M, Riedel E, Brennan MF, et al. Neo-adjuvant chemotherapy for primary high-grade extremity soft tissue sarcoma. *Ann Oncol* 2004; 15: 1667-72. <https://doi.org/10.1093/annonc/mdh431>
12. Beane JD, Yang JC, White D, Steinberg SM, Rosenberg SA, Rudloff U. Efficacy of adjuvant radiation therapy in the treatment of soft tissue sarcoma of the extremity: 20-year follow-up of a randomized prospective trial, *Ann Surg Oncol* 2014; 21: 2484-89. <https://doi.org/10.1245/s10434-014-3732-4>
13. Schmehl J, Scharpf M, Brechtel K, Kalender G, Heller S, Claussen CD, et al. Epithelioid angiosarcoma with metastatic disease after endovascular therapy of abdominal aortic aneurysm. *Cardiovasc Interv Radiol* 2011; 35: 190-3. <https://doi.org/10.1007/s00270-011-0219-6>



### OLGU SUNUMU-ÖZET

Turk J Surg 2022; 38 (3): 306-311

## Femoral arter anevrizmasından gelişen intimal anjiyosarkom: Olgu sunumu

Fatih Yanar<sup>1</sup>, Gizem Öner<sup>2</sup>, Sercan Yüksel<sup>1</sup>, Ahmet Salduz<sup>3</sup>, İbrahim Fethi Azamat<sup>1</sup>, Beyza Özçınar<sup>1</sup>, Bilge Bilgic<sup>4</sup>, Rana Günöz Cömert<sup>5</sup>, Erol Kozanoğlu<sup>6</sup>, Mehmet Kurtoğlu<sup>1</sup>

<sup>1</sup> İstanbul Üniversitesi Tıp Fakültesi, Genel Cerrahi Anabilim Dalı, İstanbul, Türkiye

<sup>2</sup> Antwerp Üniversite Hastanesi, Multidisipliner Onkolojik Merkezi (MOCA), Antwerpen, Belçika

<sup>3</sup> İstanbul Üniversitesi Tıp Fakültesi, Ortopedi ve Travmatoloji Anabilim Dalı, İstanbul, Türkiye

<sup>4</sup> İstanbul Üniversitesi Tıp Fakültesi, Patoloji Anabilim Dalı, İstanbul, Türkiye

<sup>5</sup> İstanbul Üniversitesi Tıp Fakültesi, Radyoloji Anabilim Dalı, İstanbul, Türkiye

<sup>6</sup> İstanbul Üniversitesi Tıp Fakültesi, Plastik ve Rekonstrüktif Cerrahi Anabilim Dalı, İstanbul, Türkiye

### ÖZET

İntimal anjiyosarkom nadir görülen vasküler bir malignitedir ve spesifik olmayan semptomlar nedeniyle tanı konulması oldukça zordur. İntimal anjiyosarkomların tanısı, tedavisi ve takip süreciyle ilgili tartışılmalı noktalar bulunmaktadır. Bu olgu sunumunun amacı femoral arter intimal anjiyosarkom tanısı konulan hastanın tanı ve tedavi sürecini değerlendirmek ve daha önce yapılmış olan çalışmalar doğrultusunda tartışılmalı noktaları aydınlatılmaktır. Rüptüre femoral arter anevrizma saptanarak ameliyat edilen 33 yaşında erkek hastanın patoloji sonucu intimal anjiyosarkom olarak geldi. Takiplerinde nüks gelişen hasta öncelikle kemoterapi ve radyoterapi tedavisi gördü. Tedaviye yanıt alınamaması üzerine hastaya çevre dokuları da içerecek şekilde agresif bir cerrahi yapıldı. Hastanın 10. ay kontrollerinde nüks ve metastaz saptanmadı. İntimal anjiyosarkom nadir görülse de femoral arter anevrizması saptandığında ayırıcı tanıda akla getirilmelidir. Tedavide agresif cerrahi önemlidir ancak kemo-radyoterapinin de tedaviye eklenmesi düşünülmelidir.

**Anahtar Kelimeler:** İntimal anjiyosarkom, cerrahi, kemo-radyoterapi

**DOI:** 10.47717/turksurg.2022.4381